

Criteria for Use of Levetiracetam (Keppra®)

VHA Pharmacy Benefits Management Strategic Healthcare Group and the Medical Advisory Panel

The following recommendations are based on current medical evidence and expert opinion from clinicians. The content of the document is dynamic and will be revised as new clinical data becomes available. The purpose of this document is to assist practitioners in clinical decision-making, to standardize and improve the quality of patient care, and to promote cost-effective drug prescribing. The clinician should utilize this guidance and interpret it in the clinical context of the individual patient situation.

Indication for therapy		
Patient with one of the following conditions:		
<input type="checkbox"/>	Patient diagnosed with localization-related epilepsy (producing simple or complex partial seizures with or without secondary generalization) AND ONE OF THE FOLLOWING	
<input type="checkbox"/>	Incomplete seizure control requiring adjunctive therapy OR	
<input type="checkbox"/>	Incomplete seizure control intolerant of first-line antiepileptic drugs(phenytoin, carbamazepine, divalproex sodium) OR	
<input type="checkbox"/>	Concurrent therapy with a drug that is a CYP450 substrate, inducer or inhibitor and has a narrow therapeutic range	
Dosing		
<ul style="list-style-type: none">Levetiracetam is available as 250 mg, 500 mg, and 750 mg tablets and 100mg/ml oral solution.Levetiracetam may be administered without regards to food.Initiate levetiracetam at 250 mg BID. After 2 weeks increase to 500 mg BID. If needed, additional dosing increments of 500-1000 mg/day every 2 weeks can be implemented, to a maximum recommended daily dose of 3000 mg or to maximum clinically tolerated dose.Adjust dose according to renal function, as follows:		
Creatinine Clearance(ml/min)	Dose(mg)	Interval(hours)
>80	500-1500	12
50-80	500-1000	12
30-49	250-750	12
<30	250-750	12
On dialysis	500-1000	24
(250-500 mg supplemental dose post dialysis recommended)		
Adverse Effects		
<ul style="list-style-type: none">Use of levetiracetam has been associated with neuropsychiatric adverse events, such as somnolence, fatigue, coordination difficulties, and behavioral abnormalitiesThe most common adverse events associated with levetiracetam use are asthenia (15%), somnolence (15%), headache (14%), infection (13%), dizziness (9%), and pain (7%). These adverse events appear to occur predominantly during the first 4 weeks of treatment with levetiracetam.		
Monitoring		
<ul style="list-style-type: none">Therapy with levetiracetam does not require monitoring of blood levels, white blood cell counts or liver function testsThere are no reported significant drug interactions, thus concurrent therapy with levetiracetam would not be expected to necessitate dose changes in other agents.		
Contraindications		
<input type="checkbox"/>	Hypersensitivity to levetiracetam	
Off Label Uses		
See Table 1		

References

1. Gajwani P, Forsthoff A, Muzina D, Amann B, Gao K, Elhaj O, Calabrese JR, Grunze H. Antiepileptic drugs in mood-disordered patients. *Epilepsia*. 2005;46 Suppl 4:38-44.
2. Post RM, Altshuler LL, Frye MA, Suppes T, McElroy SL, et al. Preliminary observations on the effectiveness of levetiracetam in the open adjunctive treatment of refractory bipolar disorder. *J Clin Psychiatry*. 2005 Mar;66(3):370-4.
3. Kaufman KR. Monotherapy treatment of bipolar disorder with levetiracetam. *Epilepsy Behav*. 2004 Dec;5(6):1017-20.
4. Yatham LN. Newer anticonvulsants in the treatment of bipolar disorder. *J Clin Psychiatry*. 2004;65 Suppl 10:28-35.
5. Grunze H, Langosch J, Born C, Schaub G, Walden J. Levetiracetam in the treatment of acute mania: an open add-on study with an on-off-on design. *J Clin Psychiatry*. 2003 Jul;64(7):781-4.
6. Braunig P, Kruger S. Levetiracetam in the treatment of rapid cycling bipolar disorder. *J Psychopharmacol*. 2003 Jun;17(2):239-41.
7. Goldberg JF, Burdick KE. Levetiracetam for acute mania. *Am J Psychiatry*. 2002 Jan;159(1):148.
8. Striano P, Manganelli F, Boccella P, Perretti A, Striano S. Levetiracetam in patients with cortical myoclonus: A clinical and electrophysiological study. *Mov Disord*. 2005 Aug 2; [Epub ahead of print]
9. Lim LL, Ahmed A. Limited efficacy of levetiracetam on myoclonus of different etiologies. *Parkinsonism Relat Disord*. 2005 Mar;11(2):135-7.
10. Magaudo A, Gelisse P, Genton P. Antimyoclonic effect of levetiracetam in 13 patients with Unverricht-Lundborg disease: clinical observations. *Epilepsia*. 2004 Jun;45(6):678-81.
11. Crest C, Dupont S, Leguern E, Adam C, Baulac M. Levetiracetam in progressive myoclonic epilepsy: an exploratory study in 9 patients. *Neurology*. 2004 Feb 24;62(4):640-3.
12. Recio MV, Hauser RA, Louis ED, et al. Chorea in a patient with cerebral palsy: Treatment with levetiracetam. *Movement Disorders* 2005;20(6):762-764.
13. Zesiewicz TA, Louis ED, Sullivan KL, et al. Substantial improvement in a Meige's Syndrome patient with levetiracetam treatment. *Movement Disorders* 2004;19(12):1518-1521.
14. Schauer R, Singer M, Saltuari L, Kofler M. Suppression of cortical myoclonus by levetiracetam. *Mov Disord*. 2002 Mar;17(2):411-5.
15. Frucht SJ, Louis ED, Chuang C, Fahn S. A pilot tolerability and efficacy study of levetiracetam in patients with chronic myoclonus. *Neurology*. 2001 Sep 25;57(6):1112-4.
16. Chatterjee A, Louis ED, Frucht S. Levetiracetam in the treatment of paroxysmal kinesigenic choreoathetosis. *Movement Disorders* 2002;17(3):614-615.
17. Frediani F. Anticonvulsant drugs in primary headaches prophylaxis. *Neurol Sci*. 2004 Oct;25 Suppl 3:S161-6.
18. Miller GS. Efficacy and safety of levetiracetam in pediatric migraine. *Headache*. 2004 Mar;44(3):238-43.
19. Guay DR. Oxcarbazepine, topiramate, zonisamide, and levetiracetam: potential use in neuropathic pain. *Am J Geriatr Pharmacother*. 2003 Sep;1(1):18-37.
20. Price MJ. Levetiracetam in the treatment of neuropathic pain: three case studies. *Clin J Pain*. 2004 Jan-Feb;20(1):33-6.
21. Rowbotham MC, Manville NS, Ren J. Pilot tolerability and effectiveness study of levetiracetam for postherpetic neuralgia. *Neurology*. 2003 Sep 23;61(6):866-7.
22. Wilby J, Kainth A, Hawkins N, Epstein D, McIntosh H, et al. Clinical effectiveness, tolerability and cost-effectiveness of newer drugs for epilepsy in adults: a systematic review and economic evaluation. *Health Technol Assess*. 2005 Apr;9(15):1-157, iii-iv.
23. Bergey GK. Initial treatment of epilepsy: special issues in treating the elderly. *Neurology*. 2004 Nov 23;63(10 Suppl 4):S40-8.
24. Patsalos PN. Clinical pharmacokinetics of levetiracetam. *Clin Pharmacokinet*. 2004;43(11):707-24.
25. French JA, Kanner AM, Bautista J, Abou-Khalil B, Browne T, et al. Efficacy and tolerability of the new antiepileptic drugs II: treatment of refractory epilepsy: report of the Therapeutics and Technology Assessment Subcommittee and Quality Standards Subcommittee of the American Academy of Neurology and the American Epilepsy Society. *Neurology*. 2004 Apr 27;62(8):1261-73.

27. French JA, Kanner AM, Bautista J, Abou-Khalil B, Browne T, et al. Efficacy and tolerability of the new antiepileptic drugs I: treatment of new onset epilepsy: report of the Therapeutics and Technology Assessment Subcommittee and Quality Standards Subcommittee of the American Academy of Neurology and the American Epilepsy Society. *Neurology*. 2004 Apr 27;62(8):1252-60.
28. Nash EM, Sangha KS. Levetiracetam. *Am J Health Syst Pharm*. 2001 Jul 1;58(13):1195-9.
29. Alsaadi Tm, Shatzel A, Marquez AV, et al. Clinical experience of levetiracetam monotherapy for adults with epilepsy :1 year follow up study. *Seizure* 2005;14:139-142.
30. Alsaadi TM, Koopmans S, Apperson M, Farias S. Levetiracetam monotherapy for elderly patients with epilepsy. *Seizure* 2004 Jan;13(1):58-60.
31. Cohen J. Levetiracetam monotherapy for primary generalized epilepsy. *Seizure*. 2003 Apr;12(3):150-3.
32. Ben-Menachem E, Falter U. Efficacy and tolerability of levetiracetam 3000 mg/d in patients with refractory partial seizures: a multicenter, double-blind, responder-selected study evaluating monotherapy. European Levetiracetam Study Group. *Epilepsia*. 2000 Oct;41(10):1276-83.

Table 1: Evidence for Off Label Use of Levetiracetam

Myoclonus/Dystonia	Evidence rating III-I
Lim 2005 ¹⁰	Open label trial of 8 patients over a one year period. One patient with posthypoxic myoclonus improved. The other seven did not improve or worsened.
Shauer 2002 ¹⁵	Case report, 16yo male with posthypoxic myoclonus. Improved clinical condition and allowed patient to enter rehab therapy.
Chatterjee 2002 ¹⁷	Case report, 38 yo female with paroxysmal kinesigenic choreoathetosis. Resolution of attacks however patient ceased therapy due to sedation and palpitations.
Zasiewicz 2004 ¹⁴	Case report, 56 yo female with Meige's syndrome. Minimal improvement with gabapentin, clonazepam and Botulinum toxin A. Levetiracetam 500 mg TID improved blepharospasm and oral dystonia.
Recio 2005 ¹³	Case report, 35 yo female with cerebral palsy.
Crest 2004 ¹²	Open label trial in 6 patients with progressive myoclonic epilepsy. Three patients with marked improvement, 3 patients with no improvement or worsening.
Frucht 2001 ¹⁶	Open label trial in eight patients with chronic myoclonus. Two patients discontinued therapy. Three patients experience significant reductions in their myoclonus scores.
Magaudda 2004 ¹¹	Retrospective review of 13 patients with Unverricht-Lundborg disease. Eight patients had measurable improvement on their myoclonus scores.
Bipolar Disorder	Evidence rating III-I
Kaufman 2004 ⁴	Case report. 21 yo female , failed 15 psychotropics. Patient improved over one year of followup.
Grunze 2003 ⁶	Ten patients with bipolar disease in acute mania. Used as add on therapy with haloperidol 5-10mg/day. Used an on-off design, levetiracetam till day 14, then remove and reintroduce at day 21. Seven of ten patients were responders at day 28 of the trial.
Goldberg 2002 ⁸	Case report 42 yo male. Previously treated with lithium, divalproex, carbamazepine and SSRI. Modest improvement but significant sedation and nausea which caused DC of agents. No recurrence of mania for 6 weeks on levetiracetam 2500 mg/day
Post 2005 ³	Open label adjunctive treatment of 34 patients [depression(13), mania (7) and cycling (14)]. The 16 patients with manic symptoms showed improvement in 2 weeks, four showed intervening periods of moderate to marked exacerbation.
Migraine	Evidence Rating II-2 C (for pediatric use)
Miller 2004 ¹⁹	Retrospective review of 19 pediatric patients. Eliminated headache in 10 patients, less severe headache reported in 7 patients no effect in 2 patients.
Pain	Evidence rating III C
Price 2004 ²¹	Case series of three patients, 67 yo male, 75 yo male and 55 yo female. All showed improvement on levetiracetam.
Guay 2003 ²⁰	Summary report of five case series of levetiracetam in various neuropathic pain syndromes. 48% to 60% of patients improved on therapy- outcomes were measured by visual analog scales, pain score or patient rating.
Monotherapy for epilepsy	Evidence Rating II-2 B
Alsaadi 2004 ²⁹	Retrospective review of 46 patients over a one year period. At 1 year 45% of patients were seizure free.
Alsaadi 2004 ³⁰	Retrospective review of 14 elderly patients. Eight patients became seizure free by six months
Cohen 2003 ³¹	Case report of three patients with primary generalized epilepsy. Patients became seizure free for at least 6 months on levetiracetam 1250-3000 mg/day
Ben-Menachem 2000 ³²	Randomized adjunctive therapy trial where responder patients(N=86) then underwent withdrawal to monotherapy. A responder rate of 52.9% with a mean reduction in seizure frequency by 74%

TABLE 2: Quality of Evidence [QE]

I	At least one properly done RCT
II-1	Well designed controlled trial without randomization
II-2	Well designed cohort or case-control analytic study
II-3	Multiple time series, dramatic results of uncontrolled experiment
III	Opinion of respected authorities, case reports, expert committees.

TABLE 3: Grade the Recommendation

A	A strong recommendation that the intervention is always indicated and acceptable
B	A recommendation that the intervention may be useful/effective
C	A recommendation that the intervention may be considered
D	A Recommend that a procedure may be considered not useful / effective, or may be harmful.
I	Insufficient evidence to recommend for or against – the clinician will use their clinical judgment